



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

Books

Search PubMed for

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show: 20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: J Am Coll Cardiol 2003 Mar 5;41(5):879-88

ELSEVIER SCIENCE
FULL-TEXT ARTICLE

Autologous skeletal myoblasts transplanted to ischemia-damaged myocardium in humans. Histological analysis of cell survival and differentiation.

Pagani FD, DerSimonian H, Zawadzka A, Wetzel K, Edge AS, Jacoby DB, Dinsmore JH, Wright S, Aretz TH, Eisen HJ, Aaronson KD.

Section of Cardiac Surgery, University of Michigan, Ann Arbor, MI 48109, USA. fpagani@umich.edu

OBJECTIVES: We report histological analysis of hearts from patients with end-stage heart disease who were transplanted with autologous skeletal myoblasts concurrent with left ventricular assist device (LVAD) implantation. **BACKGROUND:** Autologous skeletal myoblast transplantation is under investigation as a means to repair infarcted myocardium. To date, there is only indirect evidence to suggest survival of skeletal muscle in humans. **METHODS:** Five patients (all male; median age 60 years) with ischemic cardiomyopathy, refractory heart failure, and listed for heart transplantation underwent muscle biopsy from the quadriceps muscle. The muscle specimen was shipped to a cell isolation facility where myoblasts were isolated and grown. Patients received a transplant of 300 million cells concomitant with LVAD implantation. Four patients underwent LVAD explant after 68, 91, 141, and 191 days of LVAD support (three transplant, one LVAD death), respectively. One patient remains alive on LVAD support awaiting heart transplantation. **RESULTS:** Skeletal muscle cell survival and differentiation into mature myofibers were directly demonstrated in scarred myocardium from three of the four explanted hearts using an antibody against skeletal muscle-specific myosin heavy chain. An increase in small vessel formation was observed in one of three patients at the site of surviving myotubes, but not in adjacent tissue devoid of engrafted cells. **CONCLUSIONS:** These findings represent demonstration of autologous myoblast cell survival in human heart. The implanted skeletal myoblasts formed viable grafts in heavily scarred human

Related Articles, Links